

Is your child up to date on immunizations?

Vaccines can prevent or help protect against deadly and difficult-to-treat diseases. And the shots don't end after infancy. Check below for the recommended ages of each immunization and determine if your child is current with his or her vaccines.

Talk with your provider about an immunization schedule for your child.

Birth

- ENGERIX-B (Hepatitis B)

2 months

- Pediarix (Diphtheria, tetanus, and pertussis [DTaP]/Inactivated poliovirus/Hepatitis B)
- Prevnar 20 (Pneumococcal conjugate)
- Rotarix (Rotavirus)
- Hiberix (Haemophilus influenzae type B)

4 months

- Pediarix (Diphtheria, tetanus, and pertussis [DTaP]/Inactivated poliovirus/Hepatitis B)
- Prevnar 20 (Pneumococcal conjugate)
- Rotarix (Rotavirus)
- Hiberix (Haemophilus influenzae type B)

6 months

- Pediarix (Diphtheria, tetanus, and pertussis [DTaP]/Inactivated poliovirus/Hepatitis B)
- Prevnar 20 (Pneumococcal conjugate)
- Hiberix (Haemophilus influenzae type B)

12 months

- Priorix (Measles, Mumps, Rubella)
- Varivax (Varicella)
- Prevnar 20 (Pneumococcal conjugate)
- Havrix (Hepatitis A)

15 months

- Infanrix (Diphtheria, tetanus, and pertussis [DTaP])
- Hiberix (Haemophilus influenzae type B)*

18 months

- Havrix (Hepatitis A)

4–6 years

- ProQuad (Measles, Mumps, Rubella/Varivax [Varicella]) or separate injections (Priorix/Varivax)
- Kinrix (Diphtheria, tetanus, and pertussis [DTaP]/Inactivated poliovirus)

11–12 years

- Boostrix (Tetanus, diphtheria, and pertussis [Tdap])
- Menveo (Meningococcal Group A, C, W, Y)
- Gardasil 9 (Human papillomavirus) – 2 doses 6 months apart (over age 15 years requires 3 doses)

16 years

- Menveo (Meningococcal Group A, C, W, Y)
- Bexsero (Meningococcal Group B) – 2 doses**

Additional recommendations

Respiratory Syncytial Virus (RSV)

All infants entering into or born within their first RSV season (dependent on maternal vaccination for RSV) and all high-risk infants entering their second RSV season.

COVID-19

Children ages six months and up should get vaccinated following CDC guidance. (While the vaccine may not be available at the practice where your child is seen, the office can assist you in getting your child's vaccinations scheduled near you.)

Influenza

Yearly for all children ages 6 months and up, if no contraindications. (If it is the patient's first season of vaccination, children under 9 years should receive 2 doses separated by at least 4 weeks.)

NOTE: This schedule is for preventive vaccinations. For catch-up schedules or for patients with chronic conditions, see the CDC Immunization Schedule for vaccine recommendations.

*Can be administered as early as 12 months old

**For those ages 16-23 with no increased risk, discuss with provider.
Requires 2 doses, 6 months apart.

PED-0126-OSF-06-25



OSF[®]
HEALTHCARE